

**Research Article****Evaluation of Neuroprotective effect of medicinal plants in *Drosophila melanogaster* model****Prakash Chaudhary, Swati Dhande****HOD & Assistant Professor, Department of Pharmacology, Bharati Vidyapeeth's College of Pharmacy, Sector-8, C.B.D. Belapur, Navi Mumbai, India***ARTICLE INFO:**

Received: 25 June 2018

Received in revised form:

10 August 2018

Accepted: 20 August 2018

Available online: 30 September 2018

Keywords:*Bombax ceiba*,*Drosophila melanogaster*,

Negative,

Geotaxis assay,

Rotenone

Abstract

In this study, we investigated the neuroprotective effects of ethanolic extract of *Bombax ceiba* (EEBC) and ethanolic extract of *Gymnemasylvestre* (EEGS) against the toxicity induced by rotenone (ROT) in *Drosophila melanogaster*. Materials and Methods: Adult wild-type flies were concomitantly exposed to ROT (500 µM), EEBC (0.05% w/v and 0.1% w/v) and EEGS (0.05% w/v and 0.1% w/v) in the culture medium for 7 days. Results: ROT treated flies produced marked decreased in locomotor performance (i.e., climbing capability) in the negative geotaxis assay when compared to control group. EEBC and EEGS flies after treatment offered protection (24-42%) against the ROT-induced locomotor impairment in the negative geotaxis assay suggesting attenuation of ROT-induced locomotor deficits. Conclusion: The results of this study suggest that EEBC and EEGS were effective in reducing the toxicity induced by ROT in *D. melanogaster* as well as it confirms the significance of this model to explore possible therapeutic approaches in Parkinson's disease (PD).

Introduction

PD is a second most common neurodegenerative disease after Alzheimer's disease. It is characterized by loss of dopaminergic neurons in the substantia nigra in the mid brain resulting in depletion of dopamine, formation of filamentous intraneuronal inclusions known as Lewy bodies on the surviving dopaminergic neurons and an extrapyramidal movement disorder^[1,2]. However, free radical overproduction and abnormalities in mitochondria function have emerged as critical mediators of the dopaminergic neuronal damage^[3]. PD is clinically characterized by locomotor symptoms such as muscle rigidity, bradykinesia, postural instability and tremor^[4].

ROT a commonly used natural pesticide is a classic high affinity specific inhibitor of mitochondrial complex I which is capable of causing mitochondrial perturbations, dose-dependent ATP depletion, oxidative damage, and early mortality that mimics PD. Exposure of adult flies to sub lethal doses of ROT in the diet over 7 days causes a concentration dependent locomotor deficits, specific dopaminergic neuronal loss and reduction in the dopamine levels in adult flies^[5].

Bombax ceiba belongs to the family Bombacaceae, is one of the important medicinal plant. It contains various

phytoconstituents most common neurodegenerative diseases after

rutin, hentriacontanol, sitosterol, isomangiferin, protoatechuic acid and kaempferol etc. A paste of leaves and flowers is used as external application for skin trouble. Seeds are applied on the skin in small pox and chicken pox. Leaves are used as laxative and hematinic. Bark is used for wound healing. It has various pharmacological activities like Anti-inflammatory, Anti-diabetic, Anti-obesity, Cytotoxicity, Aphrodisiac, Hypotensive, Antioxidant, Antiangiogenic, Antimicrobial, and Antipyretic^[6].

Gymnema sylvestre belongs to the family Asclepiadaceae, popularly known as "gurmar" is a reputed herb in the Ayurvedic system of medicine used for treatment in conditions ranging from diabetes, malaria, to snakebites^[7]. The leaves contain major constituents like gymnemic acids and gymnema saponins are members of oleanane type of saponins while gymnemasides are dammarane saponins^[8,9]. Other constituents present in it are anthraquinones, hentriacontane, flavones, phytin, pentatriacontane, formic acid, butyric acid, tartaric acid, lupeol, resins, amyrin related glycosides, calcium oxalate and stigmasterol^[10]. It is used as naturopathic treatment for diabetes^[11,12]. It has also demonstrated promising effects in

the treatment of obesity, arthritis, hyperlipidemia, and hypercholesterolemia [13-15]. Furthermore, the bioactive phytochemicals have antimicrobial, antiinflammatory, and anticancer properties. The leaves are used for the treatment of obesity [16], dental caries [17], antibiotic, in stomach ache, blood purifier, and in rheumatism [18].

In the present study, the possible neuroprotective properties of EEBC and EEGS against ROT-induced oxidative stress and neurotoxicity were explored. Flies chronically exposed to ROT (500 μ M) developed selective loss of dopaminergic neurons in the brain and severe locomotor dysfunctions [19]. Dietary exposure of adult flies to EEBC (0.05% w/v and 0.1% w/v) & EEGS (0.05% w/v and 0.1% w/v) along with ROT (500 μ M) for 7 days, thereafter its ability to modulate locomotor behaviour was determined by negative geotaxis assay.

Chemicals: Rotenone (Sigma Aldrich)

Gymnemasylvestre (Family:Asclepiadaceae) leaves powder was procured from Konark Herbal and Health Care, Daman, India.

Bombaxceiba (Family:Bombaceae)bark powder wasprocured from the Rajesh chemicals Mumbai.

D.melanogaster wild-type, Canton special strain was obtained from the Indian Institute of Science and Education Research, Pune, India.

The flies were grown in length 12 cm; diameter 2 cm vials containing standard medium (8.3%, w/v maize flour; 5%, w/v glucose; 2.5%, w/v sucrose; 1.5%, w/v agar; 2%, w/v yeast powder; 0.04%, v/w propionic acid; 0.6%, v/w orthophosphoric acid; 0.7%, v/w methyl paraben) at constant temperature and humidity (23°C \pm 1°C; 60% relative humidity, respectively) under 12 h dark-light cycle. All experiments were performed with the same strain.

The powders were macerated for 8 hours using the solvent petroleum ether for defatting of the material. Crude extracts were obtained by placing 20-30 gm of powder in the soxhlet extractor using ethanol at 40°C. The crude extract that is EEBC and EEGS were further evaporated in the Rotovac evaporator and were dried to get the free flowing powder. The powders were stored in airtight container.

ROT treatedflies were exposed to low and high doses of EEBC & EEGS mixed in the culture medium. EEBC (0.05% w/v and 0.1% w/v) & EEGS (0.05% w/v and 0.1% w/v) were added in the medium at final concentrations of 5 mg/ml, 10 mg/ml. Vials of flies without test compound and ROT were used as control. Flies (male adult, 7–8 days old) were divided into six groups: (1) Control; (2) ROT (500 μ M); (3) ROT plus EEBC (0.05% w/v); (4) ROT plus EEBC (0.1% w/v); (5) ROT plus EEGS (0.05% w/v) and (6) ROT plus EEGS (0.1% w/v). The total food medium contained a volume of 1% of ROT or ROT plus EEBC or EEGS. The flies were exposed to treatments during 7 days and the vials containing flies were maintained.

The motor function was assessed using a negative geotaxis assay. During their light cycle, 15 flies were transferred into a graduated flat bottom glass tube (length 12 cm; diameter 2 cm) and allowed to habituate for at least 5 min. The tube were gently tapped at the bottom and observed for 60 sec. for the climbing activity (15 flies/trial; 3 trials were done). Locomotor behaviour was expressed as percent flies escaping beyond a minimum distance of 10 cm in 60 sec^[20].

All the data were expressed as mean \pm standard error of the mean (SEM). Statistical significance was tested using oneway ANOVA followed by Tukey's multiple comparisons test using Prism Graph Pad version 7.0 (Graph Pad software, Inc., CA). * $p < 0.05$ was considered statistically significant.

The results of this study revealed that response of the PD induced flies were significantly lower than that of control flies. Measurement of the locomotor deficits among flies of ROT treated exhibited severe locomotor impairments as evident by the large number of flies (97.73%) staying at the bottom of the glass vial. The flies co-treated with EEBC (0.05% w/v & 0.1% w/v)& EEGS (0.05% w/v & 0.1% w/v) significantly ($p < 0.05$) improved the locomotor activity when compared with ROT treated flies (Table 1 & Figure 1). The average number of flies staying at the bottom of vial was decreased to 66.63% when co-treated with EEBC & EEGS. The flies showed negative geotaxis behaviour treated with EEBC & EEGS dose dependently; indicating its neuroprotective effect.

Table1. Effect of EEBC & EEGS on locomotor activity in *Drosophila melanogaster*

Groups	No. of flies escaped beyond 10 cm distance	% protection
Vehicle Control	11.67 \pm 0.88	77.80
Disease Control	0.33 \pm 0.33	2.27
EEBC(0.05% w/v)	3.67 \pm 0.67*	24.47
EEBC(0.1% w/v)	6.34 \pm 0.33*	42.27
EEGS(0.05% w/v)	3.67 \pm 0.67*	24.47
EEGS(0.1% w/v)	6.34 \pm 0.33*	42.27

Values are expressed as mean \pm SEM; n=15, where n is the number of flies per group. *indicates there is significant difference with $p < 0.05$ when compared with disease control group by One-way ANOVA followed by Tukey's multiple comparisons test.




Fig. 1: Effect of EEBC & EEGS on locomotor activity in *Drosophila melanogaster* by negative geotaxis assay

In *Drosophila* model, ROT is commonly used for PD model. It has high affinity specific inhibitor of mitochondrial (complex I) NADH dehydrogenase. Being highly lipophilic, it crosses the blood brain barrier rapidly and accumulates in subcellular organelles like mitochondria^[21]; where oxidative damage leads to neurotoxicity of dopaminergic neurons and locomotor deficits^[22,23]. In ROT (500 μ M) induced significant locomotor impairment in flies as indicated by large number of flies remaining at bottom of vials in disease control group. The EEBC & EEGS at doses of 0.05% w/v and 0.1% w/v showed protective effect against ROT induced locomotor impairment exhibited significant improvement in locomotor activity, thus could be proved with possible action on central nervous system indicates that EETP has an ability of free radical scavenging properties helps in neuroprotection of dopaminergic neurons in brain.

In summary, this study demonstrated that EEBC & EEGS improves locomotor performance and reduce the toxicity in ROT induced *D. melanogaster* fly PD model. This study confirms the utility of this model to investigate therapeutic strategies that may be promising in the treatment of neurodegenerative diseases and can be used for studies in higher rodent model.

References

- Goedert M., Spillantini M.G., Davies S.W., Filamentous nerve cell inclusions in neurodegenerative diseases, Current Opinion in Neurology 1998;8:619-32.
- Cannon JR, Greenamyre J.T., Neurotoxic in vivo models of Parkinson's disease recent advances, Progress in Brain Research. 2010;184:17-33.
- Coulom H, Birman S. Chronic exposure to rotenone models sporadic Parkinson's disease in *Drosophila melanogaster*, The Journal of Neuroscience 2004;24: 10993-8.
- Klockgether T. Parkinson's disease: Clinical aspects, Cell and Tissue Research. 2004; 318:1:115-20.
- St. Laurent R, O'Brien L.M, Ahmad ST. Sodium butyrate improves locomotor impairment and early mortality in a rotenone induced *Drosophila* model of Parkinson's disease, Neuroscience. 2013;246:382-90.
- Verma R, Devre K, Gangrade T, Gore S, Gour SA Pharmacognostic and pharmacological overview on *Bombaxceiba*, Scholars Academic Journal of Pharmacy 2014; 3:2:100-107.
- Singh VK, Umar S, Ansari SA, Iqbal M. *Gymnemasylvestre* for diabetics, Journal of Herbs, Spices and Medicinal Plants, 2008; 14:1-2: 88-106.
- Foster S., *Gymnemasylvestre* in Alternative Medicine Reviews Monographs Thorne Research Inc, 2002; 205-207.
- Khramov VA, Spasov AA, Samokhina MP. Chemical composition of dry extracts of *Gymnemasylvestre* leaves, Pharmaceutical Chemistry Journal 2008; 42:1: 30-32.
- Sinsheimer JE, Rao GS, McIlhenny HM. Constituents from *Gymnemasylvestre* leaves. V: isolation and preliminary characterization of the gymnemic acids, Journal of Pharmaceutical Sciences 1970; 59:5:622-628.
- Shanmugasundaram ERB, Rajeswari G, Baskaran K, Kumar BRR, Shanmugasundaram KR, Ahmath BK. Use of *Gymnemasylvestre* leaf extract in the control of blood glucose in insulin-dependent diabetes mellitus, Journal of Ethno pharmacology. 1990; 30:3:281-294.
- Baskaran K, Ahmath BK, Shanmugasundaram KR, Shanmugasundaram ERB. Antidiabetic effect of a leaf extract from *Gymnemasylvestre* in non-insulin-dependent diabetes mellitus patients, Journal of Ethnopharmacology 1990; 30:3: 295-305.
- Malik JK, Manvi FV, Nanjware BR, Dwivedi DK, Purohit P, Chouhan S. Anti-arthritis activity of leaves of *Gymnemasylvestre* R.Br. leaves in rats, Der Pharmacia Lettre. 2010; 2:336-341.
- Rachh PR, Rachh MR, Ghadiya NR, Modi DC, Modi KP, Patel NM, et al. Antihyperlipidemic activity of *Gymnemasylvestre* R.Br. leaf extract on rats fed with high cholesterol diet, International Journal of Pharmacology. 2010; 6:2:138-141.

15. Spasov AA, Samokhina MP, Bulanov AE. Antidiabetic properties of *Gymnemasylvestre* (a review), Pharmaceutical Chemistry Journal. 2008; 42:11:22-26.
16. Government of India Department of Ayurveda, Yoga & Naturopathy, Unani, Siddha & Homoeopathy (AYUSH), "Mesasrngi," in The Ayurvedic Pharmacopoeia of India, The Controller of Publications, New Delhi, India, Part I, Edition 1, vol. 5, 2006:110-114.
17. Devi BP, Ramasubramaniraja R. Pharmacognostical and antimicrobial screening of *Gymnemasylvestre* R.Br, and evaluation of Gurmar herbal tooth paste and powder, composed of *Gymnemasylvestre* R.Br, extracts in dental caries, International Journal of Pharma and Bio Sciences 2010; 1:3:1-16.
18. Evans WC, Trease GE, Evans D.: Trease and Evans' Pharmacognosy. WB Saunders, Philadelphia, Pa, USA, Edition 15, 2002.
19. Hosamani R, Muralidhara. Neuroprotective efficacy of *Bacopamonnieri* against rotenone induced oxidative stress and neurotoxicity in *Drosophila melanogaster*, NeuroToxicology. 2009; 30:6:977-985.
20. Khatri DK, Juvekar AR. Abrogation of locomotor impairment in a rotenone-induced *Drosophila melanogaster* and zebrafish model of Parkinson's disease by ellagic acid and curcumin, International Journal of Nutrition, Pharmacology, Neurological Disease 2016; 6:90-6.
21. Uversky VN. Neurotoxicant induced animal models of Parkinson's disease: understanding the role of rotenone, maneb and paraquat in neurodegeneration, Cell and Tissue Research. 2004; 318:225-241.
22. Coulom H, Birman S. Chronic exposure to rotenone models sporadic Parkinson's disease in *Drosophila melanogaster*, The Journal of Neuroscience. 2004; 24: 10993-10998.
23. Hosamani R, Ramesh SR, Muralidhara. Attenuation of rotenone-induced mitochondrial oxidative damage and neurotoxicity in *Drosophila melanogaster* supplemented with creatine, Neurochemical Research 2010; 35:1402-1412

Cite this article as: **Prakash Chaudhary, Swati Dhande.** Evaluation of Neuroprotective effect of medicinal plants in *Drosophila melanogaster* model. . **Indian J. Pharm. Biol. Res.** 2018; 6(3):9-12.